



Antioxidant responses are associated with differences in drought tolerance between maize and sorghum.

Ali Ali Elnaeim Elbasheir¹ & Ludidi Ndiko^{1,2*}

¹Plant Biotechnology Research Group, Department of Biotechnology, University of the Western Cape, Bellville, South Africa

²DSI-NRF Centre of Excellence in Food Security, University of the Western Cape, Bellville, South Africa

Article info

Article history:

Received 25 April 2021

Accepted 10 June 2021

Keywords:

Reactive oxygen species, antioxidant enzyme activity, oxidative stress, maize, sorghum



Copyright©2021 JOASD

* Corresponding author

nludidi@uwc.ac.za

Conflict of Interest : The authors declare no conflict of interest

Abstract

Drought is a major cause of decreased yield in crops worldwide. Sorghum (*Sorghum bicolor*) and maize (*Zea mays*) are two of the key crops in Africa serving as human food as well as livestock feed. For improved crop production, selection for drought resilient genotypes is imperative and the biological basis for drought tolerance ought to be fully understood to achieve such selection. Sorghum can tolerate drought better than maize and it is a key model for studying the physiological and biochemical mechanisms conferring drought tolerance. In this study, comparative analyses in terms of changes in growth, chlorophyll content, ROS content, lipid peroxidation level and the activity of antioxidant enzymes were investigated. Exposure to drought triggered ROS generation in both plant species. However, sorghum showed less cell damage under water deficit compared to maize. Furthermore, differences in antioxidant enzyme activity between maize and sorghum were identified. Our findings reveal differences in and association between the physiological and biochemical responses of maize and sorghum to drought, which may be relevant for breeding drought tolerant crops.

1. INTRODUCTION

Beside the hazardous impacts on the environment, climate change is challenging socioeconomic development (He et al., 2019). Climate change and variability disturbs the basic elements of agriculture (soil moisture, heat and sunlight) as it leads to disparities in temperature, precipitation and the frequency of climate extremes such as drought (Raza et al., 2019). Drought is one of the biggest challenges affecting the globe. The lack of drought mitigation strategies and limited understanding of drought resilience, declined agricultural productivity, increases food shortages and global food insecurity (Iqbal et al., 2020). Exposure of plants to drought conditions induces oxidative

stress and reduces stomatal conductivity, which results in decreased leaf internal carbon dioxide (CO₂) and leads to over-formation of reactive oxygen species (ROS) (Xie et al., 2019). Oxidative damage via ROS generation imposes adverse changes in fundamental processes of plant growth and development, including seed germination, plant height, stem girth, leaf area, number of leaves, flowering and fruit production (Fahad et al., 2017). To counter ROS-triggered alterations, plants rely on ROS-scavenging systems which include enzymatic antioxidants such as superoxide dismutase (SOD), glutathione reductase (GR), peroxidase (POD), ascorbate peroxidase (APX), and catalase (CAT), or non-enzymatic antioxidants including carotenoids, ascorbic acid (AsA) and glutathione (GSH) (Ren

et al., 2020). The degree to which these defense systems are activated under drought stress notably varies among plant species, and even between different genotypes of the same plant species (Hasanuzzaman et al., 2020). Drought-induced oxidative stress causes lipid peroxidation as indicated by the increase in malondialdehyde (MDA) content. Thus, it is essential for plants to maintain greater levels of antioxidants in order to cope with drought stress (Laxa et al., 2019). Maize and sorghum are closely related species which are members of the Panicoideae subfamily in the family of Gramineae (Kellogg, 2001). Maize is the most important crop in the world (Li et al., 2016). In Africa, more than 300 million people depend on maize as the major staple (IITA, 1967), and is the main food crop for 50 % of the population in Sub-Saharan Africa (CGIAR, 2012). Under sufficient water supply, maize achieves high grain yield (Trout and DeJonge, 2017). However, maize is sensitive to drought stress (Meng et al., 2016). In contrast, sorghum is a drought resilient crop and the availability of its full genome sequence puts it in the lead as a key model system for deciphering the physiological and biochemical mechanisms of drought tolerance (Azzouz-Olden et al., 2020). Sorghum is the second major crop in Africa (Mundia et al., 2019). Besides being used as an animal feed, it is processed as traditional food such as semi-fermented bread, porridges, dumplings and couscous (Taylor and Belton, 2002). With the growing world population, it becomes crucial to improve and sustain crop performance under water limited conditions to ensure food security. Therefore, screening for drought adaptive responses through comparative studies in drought sensitive and drought tolerant crops is vital. It is on this basis that this study compared the physiological and antioxidant responses between maize and sorghum.

2. MATERIALS AND METHODS

2.1. Plant growth and drought treatment

Seeds of sorghum (*Sorghum bicolor* L.) Moench cv. Super Dan] and maize (*Zea mays* L.) cv. Border King] were surface sterilized and then imbibed in aerated 10 mM calcium sulphate for 16 hours prior to germination. Germinated seeds were sown in Perspex tubes (100 x 10 cm), filled with 18 L of Promix Organic (Windell Hydroponics, South Africa) saturated with water containing 1% fertilizer (v/v) [Nitrosol®, Envirogreen (Pty) Ltd)]. Seedlings were grown

until the V1 stage of development by irrigating the tubes every two days with 500 ml of water. The plants were grown in controlled greenhouse under 16/8 h day/night cycle with average temperature of 25°C. The photosynthetic photon flux density was 400 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$. At the V1 stage, control plants continued to receive 500 ml of water every second day until the day of harvest (V8 growth stage), while water-deprived plants were provided with 20% (100 ml) of that amount of water on a weekly basis until the V3 stage of development. Thereafter, the water-deficit plants were subjected to complete water deprivation until the symptoms of drought stress (leaf rolling or oldest leaves turning brown) were observed. At this point, the growth parameters of control and water-deprived plants were measured and the four youngest leaves were harvested. The harvested leaves were immediately frozen and ground into a fine powder using liquid nitrogen and then stored at -80°C. Cell viability and O₂ were determined using fresh leaves.

2.2. Trichloroacetic Acid (TCA) Extraction

A protein-free extract was obtained by homogenizing 5 volumes of 6% trichloroacetic acid (TCA) with 100 mg of powdered plant material. The homogenate was then centrifuged at 13000 x g for 15 minutes at 4°C and the supernatant was used for measuring the MDA and H₂O₂ content.

2.3. Determination of lipid peroxidation degree

Lipid peroxidation was determined by the method of Dhindsa et al. (1981). To a 200 μl aliquot of TCA extract, 400 μl of a solution containing 0.5% thiobarbituric acid (TBA) and 20% TCA was added. The mixture was boiled at 90 °C for 20 minutes and then cooled on ice. After centrifugation at 12 000 x g for 5 minutes, 200 μl of supernatant was transferred into a microtiter plate and the absorbance was recorded at 532 nm and 600 nm using a POLARstar Omega microtiter plate reader (BMG Labtech, Offenburg, Germany). The data obtained were used to calculate the MDA content based on the extinction coefficient of $1.56 \times 10^5 \text{ cm}^{-1}\text{M}^{-1}$.

2.4. Evaluation of Cell Viability

Cell viability was measured according to the method of Sanevas et al. (2007). A 1 cm² cutting

from the second youngest leaf was stained in 0.25% Evans Blue in the dark for 30 minutes at room temperature. The leaf material was then transferred into an Eppendorf tube filled with distilled water and incubated overnight at room temperature to eliminate any free dye. After the water was discarded, Evans Blue was extracted by incubating the samples in 1.5 ml of 1% sodium dodecyl sulfate (SDS) at 65°C for 1 hour. Samples were then centrifuged at 13000 x g for 5 minutes and the absorbance of the supernatant was measured at 600 nm on a POLARstar Omega microtiter plate reader (BMG Labtech, Offenburg, Germany).

2.5. Reactive oxygen species (ROS) measurements

2.5.1. Hydrogen peroxide content

A 50 µl aliquot of TCA extract was mixed with 100 µl potassium iodide (KI) and 50 µl of potassium phosphate (K₂HPO₄), pH 5.0, to a total volume of 200 µl. A standard curve with known concentrations of H₂O₂ was prepared by mixing 10 µM H₂O₂ with varying volumes of dH₂O, 0.5 M KI and 20 mM K₂HPO₄, to give a total volume of 200 µl. The absorbance was recorded at 390 nm and the extinction coefficient 39.4 mM⁻¹·cm⁻¹ was used to calculate the concentration of hydrogen peroxide in the sample.

2.5.2 Superoxide

Superoxide content was quantified according to Able et al. (1998), with modifications. A leaf disc (1 cm²) and a 4 cm cutting of the root (from the tip) collected from fresh plant tissue were incubated in a 50 mM potassium phosphate buffer (KPO₄) containing 10 mM potassium cyanide (KCN), 10 mM H₂O₂, 2% sodium dodecyl sulphate (SDS), 80 µM Thiazolyl Blue Tetrazolium Blue (MTT), for 20 minutes at room temperature. The plant tissue was then crushed with a small pestle and centrifuged at 13000 x g for 10 minutes. A 200 µl aliquot of the obtained supernatant was loaded onto a microtiter plate. Absorbance was recorded at 570 nm and the extinction coefficient 13.0 mM⁻¹·cm⁻¹ for the MTT was used to calculate super oxide content.

2.6. Total Protein Extraction and quantification

A 200 mg sample of plant tissue was homogenized in 400 µl of protein extraction buffer [40 mM phosphate buffer at pH 7.4; 1 mM

ethylenediaminetetraacetic acid (EDTA) and 5% (w/v) polyvinylpolypyrrolidone (PVPP)]. The homogenate was centrifuged at 13000 x g for 20 minutes at 4°C and the supernatant was used as enzyme source for APX, CAT, SOD and GR assays. The concentration of protein in the extracts was estimated by the method of Bradford (1976) using bovine serum albumin (BSA) as a standard.

2.7. Assays of antioxidant enzyme activities

For native in-gel assays, the staining of all enzymes was done using non-denaturing discontinuous gel electrophoresis formed of 15 % separating gel [except for CAT (in which case the separating gel was a 7.5 % gel)] and 5 % stacking gel. The gels were electrophoresed with a running buffer containing 25 mM Tris, 192 mM glycine, and additionally (in the case of APX) 2 mM ascorbic acid (AsA) at 4°C and 80 mV. All gels were run until the loading dye reached the bottom edge of the gel except for CAT (the gel was allowed to run for 2 hours after the dye ran off the gel to allow for good separation of isoforms).

2.7.1. Assay of APX activity

Following electrophoretic separation, the ascorbate peroxidase gels were stained according to Seckin et al. (2010) in order to determine the activity of each of the APX isoforms in maize and sorghum. The ascorbate peroxidase total activity was determined by the method of Nakano and Asada (1981). APX activity was estimated by measuring the decrease in absorbance at 290 nm as ascorbate was oxidised and using the extinction coefficient of 2.8 mM⁻¹·cm⁻¹ for ascorbate. One unit of APX activity was defined as the amount of protein required to hydrolyse 1 µmole of H₂O₂/minute.

2.7.2. Assay of CAT activity

The staining of catalase gels was performed as described by Yamashita et al. (2007). The catalase total activity was measured by following the consumption of hydrogen peroxide as described by Lück (1965). The calculations were done by monitoring the decrease in absorbance at 240 nm ($\epsilon = 39.4 \text{ mM}^{-1} \text{ cm}^{-1}$), one unit of enzyme is the amount necessary to convert 1 µmole of H₂O₂/ minute.

2.7.3. Assay of SOD activity

Superoxide dismutase (SOD) activity staining was assayed according to Beauchamp and Fridovich (1971). One unit of SOD activity represented the amount of enzyme required to inhibit 50 % of NBT.

2.7.4. Assay of GR activity

Following native PAGE separation, Glutathione reductase (GR) activity staining was done as reported by Rao et al. (1996). The total activity of GR was estimated by the method of Foyer and Halliwell (1976). GR activity was expressed as μmol of NADPH oxidized/minute per mg of protein.

2.8 Statistical analysis

Data generated were subjected to a one-way analysis of variance (ANOVA) and significance was determined by the Tukey-Kramer test at a 5% level of significance using GraphPad Prism 6.0.

3. RESULTS AND DISCUSSION

3.1. Reduction in growth under drought is more severe in maize than in sorghum

Exposure of crops to drought can adversely affect their growth and yield (Hussain et al., 2019). Under drought stress, maize suffered a greater reduction of shoot length (approximately

29 %) than sorghum which showed only 16 % decrease in shoot length (Fig. 1. a). Both maize and sorghum exhibited a significant increase in their root length under drought as shown in Fig. 1. b. However, the increase in root length was even greater in sorghum (approximately 44 %) than in maize (approximately 14 %), similarly to the study reported by Wright et al. (1983). Drought decreased the shoot fresh weight in both maize and sorghum by approximately 84 % and 77 % respectively (Fig.1. c). The longer root length in sorghum is likely an adaptive response in order to reach deeper soil layers to acquire moisture.

3.2. Oxidative stress is higher in maize than sorghum under water deprivation

Fig. 2 illustrates hydrogen peroxide content, superoxide content and the level of lipid peroxidation (MDA content) in maize and sorghum. Drought stress increased the accumulation of H_2O_2 in the leaves of maize and sorghum, while no marked changes in H_2O_2 content were observed in the roots of both species (Fig. 2. a and b). However, maize plants showed a much higher increase in H_2O_2 content, as it increased by approximately 70%, than the approximately 55% increase in H_2O_2 content observed in sorghum. Under drought stress, $\text{O}_2^{\cdot -}$ content was increased by approximately 38.6%




Fig. 1. The effect of drought stress on shoot length (a), root length (b) and shoot fresh weight (c) in *Zea mays* and *Sorghum bicolor*. The stress parameters of well-watered and water-deprived plants were determined at V8 stage of growth. Values are means \pm SE of ten plants, $P \leq 0.05$.




Fig. 2. The degree of oxidative stress exhibited in *Zea mays* and *Sorghum bicolor* under well-watered and water-deprived conditions. Hydrogen peroxide (H₂O₂) content in leaves (a) and roots (b), and O₂⁻ in leaves (c) and roots (d) were measured as the indication of ROS accumulation; and lipid peroxidation in leaves (e) and roots (f) was measured as an indication of oxidative stress. Data presented are means (±SE) of three independent experiments (n=3). Different letters above error bars denote mean values that are significantly different at P ≤ 0.05.

in the leaves of maize, whereas no significant change in O₂⁻ content was observed in sorghum (Fig. 2. c). Similarly, an increase of approximately 25% in O₂⁻ content in the roots occurred in water-deprived maize, whereas no significant difference in O₂⁻ was observed in sorghum (Fig. 2. d). Drought increased MDA content by approximately 58% in maize leaves

and by approximately 41% in sorghum leaves (Fig. 2. e). In the roots (Fig. 2. f), maize plants showed a 70% increase in MDA content in response to drought while sorghum plants showed a 55% increase in MDA content. The above results show that maize suffers greater oxidative damage than sorghum under the same drought conditions. The drought-




Fig. 3. The effect of drought stress on cell viability in *Zea mays* and *Sorghum bicolor*. Data presented are means (\pm SE) of three independent experiments ($n=3$). Different letters above error bars denote mean values that are significantly different at $P \leq 0.05$.

induced accumulation of H_2O_2 and $O_2^{\cdot-}$ content demonstrated here are the cause of the higher lipid peroxidation and the differences in their accumulation between maize and sorghum partly explain the differing degree of tolerance to drought stress between these two species.

Oxidative stress occurs when the critical balance between ROS formation and ROS scavenging by antioxidant enzymes is disturbed due to decreased antioxidant enzyme activity or excess formation of ROS, or both (Poljsak et al., 2013).

This excessive ROS accumulation is caused by a wide range of environmental stresses, including drought (Cruz de Carvalho, 2008).

3.3. Drought stress induces cell death in maize and sorghum

Drought stress affects photosynthesis and enhances photorespiration, disrupts normal cell homeostasis and leads to increased formation of reactive oxygen species (ROS) (Miller et al., 2010). The excess ROS may cause severe oxidative stress and ultimately cell death (Sharma et al., 2012). However, an effective ROS scavenging system and low levels of cell death under drought conditions are associated with plant tolerance to drought stress (Laxa et al., 2019).

Fig. 3. shows the level of cell death as determined by Evans Blue uptake. Under drought, maize showed nearly 12 % increase in the level of cell death, while cell death in sorghum was approximately 7 % higher when compared to their relative controls. The more pronounced increase of cell death in maize could be attributed to the higher oxidative stress in maize compared to sorghum as a result of excessive ROS accumulation under drought conditions (Fig. 2. e and f), similar findings were described for salinity stress by Keyster et al. (2013).

3.4 Drought stress significantly alters the activity of antioxidant enzymes

During ROS-triggered oxidative stress, plants regulate their antioxidant defense system to scavenge ROS, thus avoiding excessive ROS accumulation and contributing to plant survival




Fig. 4. Changes in ascorbate peroxidase (APX) isoforms in response to drought stress in the leaves (a) and roots (b) of *Zea mays* and *Sorghum bicolor*. WW represents well-watered, WD represents water-deprived. White bands in dark-blue background signify isoforms of APX.

(Huang et al., 2019). Based on APX native PAGE activity staining, four isoforms were detected in maize while only three isoforms were identified in sorghum (Fig. 4.). As a result of exposure to drought, ascorbate peroxidase isoform activity was differentially regulated between maize and sorghum. Interestingly, drought increased APX total activity by approximately 38% in the leaves of maize and by approximately 19% in the leaves of sorghum when compared to their respective controls (Fig. 5. a). However, in response to drought, APX total activity was decreased by approximately 73% in the roots of maize, whereas it remained unchanged in sorghum (Fig. 5. b). The increase in APX activity could be

According to the native PAGE analysis, three CAT isozymes were detected in the leaves of maize and sorghum. In the roots, only one CAT isoform was identified in maize, while two CAT isoforms were detected in sorghum. The activity of all CAT isoforms was significantly attenuated in both maize and sorghum in response to drought stress (Fig. 6.). In response to drought, the total CAT activity was decreased by approximately 45% in the leaves of maize and by approximately 33% in the leaves of sorghum (Fig. 7. a). Similarly, an approximately 41% and 66% reduction in CAT activity was observed in the roots of maize and sorghum plants, respectively, under water deprivation (Fig. 7. b).




Fig. 5. Determination of the total APX activity in leaves (a) and roots (b) of *Zea mays* and *Sorghum bicolor* by spectrophotometry. Error bars represent the means \pm SE; n= 3. Different letters indicate statistically




Fig. 6. Catalase isoforms activity was decreased under water stress in leaves (a) and roots (b) of *Zea mays* and *Sorghum bicolor*. WW represents tissue obtained from well-watered plants, WD represents tissue obtained from water-deprived plants.

attributed to efforts to avoid oxidative damage caused by the excessive generation of H₂O₂. Similar findings were reported by Chugh et al. (2011), who observed a significant increase in APX activity in maize when exposed to water deficit stress. The greater increase of APX activity in the leaves of maize could be a result of the higher increase of H₂O₂ content in maize than in sorghum under drought.

The results obtained above are in agreement with a study done by Bakalova et al. (2004) in wheat and Pan et al. (2006) in liquorice. Decreased catalase activity could be the reason for the observed increase in H₂O₂ content in response to drought.




Fig. 7. Determination of total catalase activity in leaves (a) and roots (b) of *Zea mays* and *Sorghum bicolor* by a spectrophotometry assay. Error bars represent the means \pm SE; n = 3. Different letters indicate statistically different means ($P \leq 0.05$).




Fig. 8. Changes in superoxide dismutase (SOD) isoenzyme activities in native gels in leaves and roots of *Zea mays* and *Sorghum bicolor* in response to drought stress. The three in-gel assays represent the detection of different SOD isoforms (a) without inhibitors, (b) treated with 5 mM KCN and (c) treated with 6 mM H_2O_2 . WW represents tissue from well-watered plants, WD represents tissue from water-deprived plants, L represents leaves and R represents the roots.




Fig. 9. Spectrophotometric determination of total SOD activity in the leaves (a) and roots (b) of *Zea mays* and *Sorghum bicolor*. Error bars represent the means \pm SE; n = 3. Different letters indicate significant differences between means at $P \leq 0.05$.

Based on the sensitivity to selective SOD inhibitors (5 mM KCN and 6 mM H₂O₂), a total of six SOD isozymes were identified in maize and sorghum (Fig. 8.). Two isoforms were identified as manganese superoxide dismutase (MnSOD), three as copper/zinc superoxide dismutase (Cu/Zn SOD) and one as iron superoxide dismutase (Fe-SOD). As shown in Fig. 9. a, overall SOD activity slightly increased (by approximately 12%) in the leaves of water-deprived maize plants, while no significant change was observed in the leaves of water-deprived sorghum plants. The upregulation of SOD activity in maize is likely a response to the higher O₂⁻ accumulation in maize than in sorghum in response to drought, similarly to work described by Luna et al. (1985), where higher SOD activity was observed in maize than

wheat when subjected to water deprivation. Drought decreased total SOD activity in the roots of both maize and sorghum (by 20% and 22%, respectively) (Fig. 9. b). The downregulation of SOD activity in maize roots is likely the underlying reason for the elevation of O₂⁻ content.

Although the SOD activity was also inhibited in sorghum roots by drought, no significant difference in O₂⁻ content was observed, suggesting that sorghum has superior SOD capacity than maize.

Glutathione reductase enzyme (GR) is essential for restoration of the reduced glutathione (GSH) pool by converting the oxidised glutathione disulphide (GSSG) back into the sulfhydryl form [GSH donates an electron to dehydroascorbate reductase (DHAR), which




Fig. 10. Changes in glutathione reductase (GR) isozyme activity in native gels in leaves and roots of *Zea mays* and *Sorghum bicolor* in response to drought stress. WW represents well-watered plants, WD represents water-deprived plants, L represents the leaves and R represents the roots.




Fig. 11. Spectrophotometric determination of total GR activity in the leaves (a) and roots (b) of *Zea mays* and *Sorghum bicolor*. Error bars represent the means ± SE; n = 3. Different letters indicate significant differences between means at P ≤ 0.05.

reduces dehydroascorbic acid (DHA to ascorbate] (Meloni et al., 2003). Based on native PAGE gel analysis, a total of six GR isozymes are present in maize and sorghum. Drought stress significantly altered the glutathione reductase isoform activity between maize and sorghum as shown in Fig. 10. The overall activity of GR increased by approximately 33% in the leaves of maize under drought stress, whereas no statistically significant difference was observed in sorghum in response to drought (Fig. 11. a). The increase in GR activity observed in the leaves of maize is likely due to the upregulation of APX activity in the leaves of maize as the cells may be driving GR activity for reformation of both GSH and ascorbate. GR total activity in the roots of maize was decreased by approximately 16% in response to drought, while the GR activity was significantly higher (by approximately 22%) in sorghum roots (Fig. 11. b). The reduction in GR activity in the roots of maize could similarly be attributed to the downregulation of APX activity in the roots of maize as the cells may have less need for driving GR activity for reformation of both GSH and ascorbate.

4. CONCLUSION

Drought stress significantly inhibited the growth of maize and induced ROS, lipid peroxidation and cell death. In contrary, sorghum exhibited less effects on growth and oxidative stress in response to drought, which may be due to differing levels of antioxidant enzyme activity in these plant species. From our results, it appears that adaptive responses in enzymatic antioxidants in maize were less sufficient than in sorghum to prevent oxidative damage caused by drought stress. This may provide the foundation for further elucidating the differences in molecular events that determine differences in drought tolerance between maize and sorghum, likely through omics-based analyses that could assist in improvement of maize drought tolerance. This can be achieved if regulation of the expression of genes discovered to be key to sorghum tolerance can be manipulated in maize to emulate the expression profile of sorghum.

ACKNOWLEDGMENTS

This work was supported by the National Research Foundation (South Africa), the Department of Science and Technology/National Research Foundation Centre of Excellence in

Food Security and the University of the Western Cape.

REFERENCES

- Able, A.J., Guest, D.I., Sutherland, M.W. (1998). Use of a new tetrazolium-based assay to study the production of superoxide radicals by tobacco cell cultures challenged with a virulent zoospores of *Phytophthora parasitica varnicotianae*. *Plant Physiology* 117, 491-499.
- Azzouz-Olden, F., Hunt, A.G. Dinkins, R. (2020). Transcriptome analysis of drought-tolerant sorghum genotype SC56 in response to water stress reveals an oxidative stress defense strategy. *Molecular Biology Reports* 47, 3291-3303.
- Bakalova, S., Nikolova, A., Nedeva, D. (2004). Isoenzyme profiles of peroxidase, catalase and superoxide dismutase as affected by dehydration stress and ABA during germination of wheat seeds. *Bulgarian Journal of Plant Physiology* 30, 64.
- Beauchamp, C., Fridovich, I. (1971). Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Analytical Biochemistry* 44, 276-287.
- CGIAR research program on maize (CRP) (2012). Why maize. <https://maize.org/why-maize/> (accessed 08.04.2021).
- Chugh, V., Kaur, N., Gupta, A.K. (2011). Evaluation of oxidative stress tolerance in maize (*Zea mays* L.) seedlings in response to drought. *Indian Journal of Biochemistry & Biophysics* 48, 47-53.
- Cruz de Carvalho, M.H. (2008). Drought stress and reactive oxygen species: production, scavenging and signaling. *Plant Signaling & Behavior* 3, 156-165.
- Dhindsa, R.S., Plumb-Dhindsa, P.A.M.E.L.A., Thorpe, T.A. (1981). Leaf senescence: correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *Journal of Experimental Botany* 32, 93-101.
- Fahad, S., Bajwa, A.A., Nazir, U., Anjum, S.A., Farooq, A., Zohaib, A., Sadia, S., Nasim, W., Adkins, S., Saud, S., Ihsan, M.Z. (2017). Crop production under drought and heat stress: plant responses and management options. *Frontiers in Plant Science* 8, 1147.
- Foyer, C.H., Halliwell, B. (1976). The presence of glutathione and glutathione reductase in

- chloroplasts: a proposed role in ascorbic acid metabolism. *Planta* 133, 21-25.
- Hasanuzzaman, M., Bhuyan, M.H.M., Zulfiqar, F., Raza, A., Mohsin, S.M., Mahmud, J.A., Fujita, M., Fotopoulos, V. (2020). Reactive oxygen species and antioxidant defense in plants under abiotic stress: revisiting the crucial role of a universal defense regulator. *Antioxidants* 9, 681.
- He, X., Estes, L., Konar, M., Tian, D., Anghileri, D., Baylis, K., Evans, T.P., Sheffield, J. (2019). Integrated approaches to understanding and reducing drought impact on food security across scales. *Current Opinion in Environmental Sustainability* 40, 43-54.
- Huang, H., Ullah, F., Zhou, D.X., Yi, M., Zhao, Y. (2019). Mechanisms of ROS regulation of plant development and stress responses. *Frontiers in Plant Science* 10, 800.
- Hussain, H.A., Men, S., Hussain, S., Chen, Y., Ali, S., Zhang, S., Zhang, K., Li, Y., Xu, Q., Liao, C., Wang, L. (2019). Interactive effects of drought and heat stresses on morpho-physiological attributes, yield, nutrient uptake and oxidative status in maize hybrids. *Scientific Reports* 9, 1-12.
- Iqbal, M.S., Singh, A.K., Ansari, M.I. (2020). Effect of drought stress on crop production, in Rakshit, A., Singh, H., Singh, A.K., Singh, U.S., Fraceto, L. (Eds.), *New Frontiers in Stress Management for Durable Agriculture*. Springer, Singapore, pp.35-47.
- Kellogg, E.A. (2001). Evolutionary history of the grasses. *Plant Physiology* 125, 1198-1205.
- Keyster, M., Klein, A., Du Plessis, M., Jacobs, A., Kappo, A., Kocsy, G., Galiba, G., Ludidi, N. (2013). Capacity to control oxidative stress-induced caspase-like activity determines the level of tolerance to salt stress in two contrasting maize genotypes. *Acta Physiologiae Plantarum* 35, 31-40.
- Laxa, M., Liebthal, M., Telman, W., Chibani, K., Dietz, K.J. (2019). The role of the plant antioxidant system in drought tolerance. *Antioxidants* 8, 94.
- Li, Y., Bingxin, Y., Dongxing, Z., Tianliang, Z., Yunxia, W., Tao, C. (2016). Research progress on precision planting technology of maize. *Transactions of the Chinese Society for Agricultural Machinery* 47, 38-48.
- Lück, H., (1965). Catalase, in: Bergmeyer, H-U. (Eds.), *Methods of Enzymatic Analysis*. Academic Press, Cambridge, Massachusetts, pp. 885-894.
- Luna, M., Badiani, M., Felici, M., Artemi, F., Sermanni, G.G. (1985). Selective enzyme inactivation under water stress in maize (*Zea mays* L.) and wheat (*Triticum aestivum* L.) seedlings. *Environmental and Experimental Botany* 25, 153-156.
- Meloni, D.A., Oliva, M.A., Martinez, C.A., Cambraia, J. (2003). Photosynthesis and activity of superoxide dismutase, peroxidase and glutathione reductase in cotton under salt stress. *Environmental and Experimental Botany* 49, 69-76.
- Meng, Q., Chen, X., Lobell, D.B., Cui, Z., Zhang, Y., Yang, H., Zhang, F. (2016). Growing sensitivity of maize to water scarcity under climate change. *Scientific Reports* 6, 1-7.
- Miller, G.A.D., Suzuki, N., Ciftci-Yilmaz, S.U.L.T.A.N., Mittler, R.O.N. (2010). Reactive oxygen species homeostasis and signalling during drought and salinity stresses. *Plant, Cell & Environment* 33, 453-467.
- Mundia, C.W., Secchi, S., Akamani, K., Wang, G. (2019). A regional comparison of factors affecting global sorghum production: The case of North America, Asia and Africa's Sahel. *Sustainability* 11, 2135.
- Nakano, Y., Asada, K. (1981). Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant and Cell Physiology* 22, 867-880.
- Pan, Y., Wu, L.J., Yu, Z.L. (2006). Effect of salt and drought stress on antioxidant enzymes activities and SOD isoenzymes of liquorice (*Glycyrrhiza uralensis* Fisch). *Plant Growth Regulation* 49, 157-165.
- Poljsak, B., Šuput, D., Milisav, I. (2013). Achieving the balance between ROS and antioxidants: when to use the synthetic antioxidants. *Oxidative Medicine and Cellular Longevity* 2013, 956792.
- Rao, M.V., Paliyath, G., Ormrod, D.P. (1996). Ultraviolet-B-and ozone-induced biochemical changes in antioxidant enzymes of *Arabidopsis thaliana*. *Plant Physiology* 110, 125-136.
- Raza, A., Razzaq, A., Mehmood, S.S., Zou, X., Zhang, X., Lv, Y., Xu, J. (2019). Impact of climate change on crops adaptation and strategies to tackle its outcome: A review. *Plants* 8, 34.
- Ren, R., Li, Z., Zhang, L., Zhou, H., Jiang, X., Liu, Y. (2020). Enzymatic and nonenzymatic antioxidant systems impact the viability of cryopreserved *Paeonia suffruticosa* pollen. *Plant Cell, Tissue and Organ Culture (PCTOC)* 144, 233-246.
- Sanevas, N., Sunohara, Y., Matsumoto, H. (2007). Characterization of reactive oxygen species-involved oxidative damage in *Hapalosiphon* species crude extract-treated

- wheat and onion roots. *Weed Biology and Management* 7, 172-177.
- Seckin, B., Turkan, I., Sekmen, A.H., Ozfidan, C. (2010). The role of antioxidant defense systems at differential salt tolerance of *Hordeum marinum* Huds (sea barley grass) and *Hordeum vulgare* L. (cultivated barley). *Environmental and Experimental Botany* 69, 76-85.
- Sharma, P., Jha, A.B., Dubey, R.S., Pessaraki, M. (2012). Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Journal of*
- Taylor J.R.N., Belton P.S. (2002). Sorghum, in: Belton, P.S., Taylor, J.R.N. (Eds), *Pseudocereals and Less Common Cereals*. Springer, Berlin, Heidelberg, pp. 25-91.
- The International Institute of Tropical Agriculture (IITA) (1967). Maize. <https://www.iita.org/cropsnew/maize/> (accessed 08.04.2021).
- Trout, T.J., DeJonge, K.C. (2017). Water productivity of maize in the US high plains. *Irrigation Science* 35, 251-266.
- Wright, G.C., Smith, R.C.G., Morgan, J.M. (1983). Differences between two grain sorghum genotypes in adaptation to drought stress. III. Physiological responses. *Australian Journal of Agricultural Research* 34, 637-651.
- Xie, X., He, Z., Chen, N., Tang, Z., Wang, Q., Cai, Y. (2019). The roles of environmental factors in regulation of oxidative stress in plant. *BioMed Research International* 2019, 9732325.
- Yamashita, K., Shiozawa, A., Banno, S., Fukumori, F., Ichiishi, A., Kimura, M., Fujimura, M. (2007). Involvement of OS-2 MAP kinase in regulation of the large-subunit catalases CAT-1 and CAT-3 in *Neurospora crassa*. *Genes & Genetic Systems* 82, 301-310.